

Streptokinase clot culture compared with whole blood culture for
isolation of Salmonella typhi and S. paratyphi A from
patients with enteric fever

2

DTIC FILE COPY

C.H. Simajuntak, S.L. Hoffman, R. Darmowigoto, M. Lesmana,
Soeprawoto and D.C. Edman

REPORT NO.

TR - 1046

AD-A201 889



**UNITED STATES NAVAL
MEDICAL RESEARCH UNIT NO. TWO**

APO SAN FRANCISCO, CALIFORNIA 96328

NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND
BETHESDA, MARYLAND

88 1031 114

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Distribution of this document is unlimited.		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NAMRU-2-TR-1046			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION U.S. Naval Medical Research Unit No. 2		6b. OFFICE SYMBOL (if applicable) NAMRU-2	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) AFPO San Francisco, California 96528-5000			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Naval Medical Research & Development Command		8b. OFFICE SYMBOL (if applicable) NMRDC	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code) Bethesda, Maryland 20814			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 61102A	PROJECT NO. M161102BS10	TASK NO. AF
					WORK UNIT ACCESSION NO. 428
11. TITLE (Include Security Classification) Streptokinase clot culture compared with whole blood culture for isolation of <u>Salmonella typhi</u> and <u>S. paratyphi</u> A from patients with enteric fever.					
12. PERSONAL AUTHOR(S) C.H. Simanjuntak, S.L. Hoffman, R. Darmowigoto, M. Lesmana, Soeprawoto & D.C. Edman, NAMRU-2 and Indonesian Department of Health.					
13a. TYPE OF REPORT Technical Report		13b. TIME COVERED FROM 1984 TO 1985		14. DATE OF REPORT (Year, Month, Day) 1988	
				15. PAGE COUNT 2	
16. SUPPLEMENTARY NOTATION Published in the Trans. R. Soc. Trop. Med. Hyg., 82:340-341. This study was done in cooperation with the Indonesian Ministry of Health.					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
			Streptokinase clot culture <u>S. paratyphi</u>		
			<u>Salmonella typhi</u> enteric fever		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>→ The sensitivities of whole blood and clot cultures were compared in 155 patients with typhoid or paratyphoid fever. <u>Salmonella typhi</u> or <u>S. paratyphi</u> A were isolated from 98.7% of 5 ml 1:10 blood:broth ratio blood cultures and 94.8% of 5 ml streptokinase clot cultures ($P > 0.05$). There was no difference in speed of isolation. Whole blood culture and clot culture were of nearly equal sensitivity in this group of patients. <i>Reprints sent to</i></p>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL RAM I: Groves, M.			22b. TELEPHONE (Include Area Code) 301/663-7567		22c. OFFICE SYMBOL NMRDC

DD Form 1473, JUN 76

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

S/N 0102-LF-014-6603

UNCLASSIFIED

ADMINISTRATIVE INFORMATION

C.G. HAYES, Ph.D.
Scientific Director

F.P. PALEOLOGO
Officer in Charge
Jakarta Detachment

This study was supported by the
World Health Organization fund No.83141,
T6/181/25; Ministry of Health, Indonesia;
Pertamina Oil Co., Jakarta, Indonesia;
and the Naval Medical Research and
Development Command, Navy Department for
Work Unit No. 3M161102BS10.AF428.

Distribution of this document is
unlimited.

J.C. COOLBAUGH
CAPT MSC USN
Commanding Officer



Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	20

Streptokinase clot culture compared with whole blood culture for isolation of *Salmonella typhi* and *S. paratyphi* A from patients with enteric fever

Cyrus H. Simanjuntak, Stephen L. Hoffman*, Ruwido Darmowigoto, Murad Lesmana, Soeprawoto and David C. Edman National Institute of Health Research and Development, Ministry of Health, Jakarta, Indonesia; United States Naval Medical Research Unit No. 2, Jakarta Detachment, Jakarta, Indonesia; and the Department of Health, Pertamina Oil Company, Plaju and Jakarta, Indonesia.

Abstract

The sensitivities of whole blood and clot cultures were compared in 155 patients with typhoid or paratyphoid fever. *Salmonella typhi* or *S. paratyphi* A were isolated from 98.7% of 5 ml 1:10 blood:broth ratio blood cultures and 94.8% of 5 ml streptokinase clot cultures ($P>0.05$). There was no difference in the speed of isolation. Whole blood culture and clot culture were of nearly equal sensitivity in this group of patients.

Introduction

Confirmation of the diagnosis of typhoid and paratyphoid fever requires isolation of the causative organism from a patient. The bone marrow aspirate culture is the single most sensitive method for isolating *Salmonella typhi* and *S. paratyphi* A from patients with enteric fever (HOFFMAN *et al.*, 1984, 1986; GILMAN *et al.*, 1975; GUERRA-CACERAS *et al.*, 1979; BENAVENTE *et al.*, 1984). However, obtaining bone marrow aspirates is not always possible. WATSON (1978) reported that the streptokinase clot culture (STKCC) was 28% more sensitive than whole blood culture (BC), a finding that was not confirmed in Jakarta (HOFFMAN *et al.*, 1986). This study was designed to compare the sensitivities of STKCC and whole BC for isolating *S. typhi* and *S. paratyphi* A in another location.

Methods

Patient selection

From 27 July 1984 to 8 April 1985, all patients presenting to the inpatient and outpatient departments of Pertamina Oil Company Hospital in Plaju, Sumatera, Indonesia were intended to have 5 ml STKCCs and 5 ml BCs. In some cases specimens for both cultures were not obtained. Only patients from whom specimens for both cultures were obtained, and

who had at least one specimen positive for *S. typhi* or *S. paratyphi* A, are included in this report.

Acquisition and processing of specimens

5 ml of blood were placed in 45 ml of 10% Oxgall (Difco Laboratories, Detroit, Michigan, USA) at the bedside (5 ml 1:10 BC) and 5 ml of blood were placed in a sterile tube and allowed to clot at room temperature. Within 24 h (90% within 3 h) the blood was centrifuged for 15 min at 1000 g. The serum was removed and the clot was placed in 15 ml of 10% Oxgall with 1500 units of streptokinase (Kabikinase, Pharmacia Laboratories, Piscataway, New Jersey, USA) (5 ml STKCC).

Isolation and biochemical and serological identification of cultured organisms were done according to standard methods described by EDWARDS & EWING (1972). Aliquots from the Oxgall were inoculated on to MacConkey's and salmonella-shigella agar media (Difco) daily for several days or until colonies resembling *Salmonella* were cultured. Cultures that did not yield an isolate after 7 d were considered negative and discarded. Colonies resembling *Salmonella* were tested with *Salmonella* group-specific antisera (Bio-Merieux, 69260 Charbonnières les Bains, France). All isolates were sent to Jakarta for confirmation at the National Institute of Health Research and Development and US NAMRU-2 laboratories.

Results

Patient population

155 patients had at least one of the cultures positive for *S. typhi* (93) or *S. paratyphi* A (62). There were 83 males and 72 females with an age of 12.9 ± 8.7 (1-43) years [mean \pm SD, (range)]. They had been ill for 5.8 ± 3.3 (2-15) d before specimen acquisition and 7.8% had taken an antibiotic before blood was drawn.

Isolation rates

Salmonella sp. was isolated from 5 ml 1:10 BC in 98.7% of cases and from STKCC in 94.8% of cases

Table—Isolation rates of *Salmonella typhi* and *S. paratyphi* A using blood culture and STKCC among the 155 patients who had at least one positive culture.

Organism	No.	BC ¹ positive No.	%	STKCC ¹ positive No.	%
<i>S. typhi</i>	93	92	98.9	89	95.7
<i>S. paratyphi</i> A	62	61	98.4	58	93.5
<i>S. typhi</i> or <i>S. paratyphi</i> A	155	153	98.7	147	94.8

¹BC = 5 ml 1:10 blood:broth ratio whole blood culture, and STKCC = 5 ml streptokinase clot culture.

*Present address: Infectious Diseases Department, Naval Medical Research Institute, Bethesda, MD20814-5055, USA.

The opinions and assertions contained herein are those of the authors and are not to be construed as official or reflecting the views of the Navy department or the naval service at large or the Indonesian Ministry of Health.

Official reprint requests should be sent to: Publications Office, U.S. Naval Medical Research Unit No. 2, APO San Francisco, CA 96328, USA or Accelerando Building, 395 Buendia Extension, Makati, Metro Manila, Philippines.

Please address all correspondence to: Dr Cyrus H. Simanjuntak, National Institute of Health Research and Development, Ministry of Health, Jalan Percetakan Negara 29, Jakarta, Indonesia.

($P > 0.05$, McNemar's test) (Table 1). The BC became positive in 1.78 ± 0.37 and 1.65 ± 0.87 d after acquisition and the STKCC in 1.82 ± 0.86 and 1.65 ± 0.83 d for *S. typhi* and *S. paratyphi* A respectively.

Discussion

The 5 ml 1:10 BC and STKCC were of nearly equal sensitivity for isolating *S. typhi* and *S. paratyphi* A from patients with typhoid and paratyphoid fever. In Jakarta the 8 ml 1:10 BC was 11% more sensitive than the STKCC. The results of both studies in Indonesia differ markedly from those of WATSON (1978), who showed that the 8 ml STKCC was 28% more sensitive than 8 ml 1:7.5 BC. Watson used bile salt broth and streptokinase/streptodornase (Varidase, Lederle) and we used 10% Oxgall and streptokinase. A series of unpublished experiments in Jakarta has shown that there is no advantage of bile salt broth over Oxgall (M. Lesmana, personal communication). We doubt whether Varidase, which is no longer available, would have been superior to streptokinase. It is possible, but unlikely, that differences in patient populations such as age, pre-culture antibiotic exposure, and pre-culture length of illness could account for the difference.

The STKCC offers the advantage of providing material for culture and serum for serological studies and the disadvantage of being more difficult to process, more expensive and perhaps slightly less sensitive than whole blood culture. When the only source of specimens is clots remaining from blood collected for serological studies, STKCC is an adequate substitute for whole BC.

Acknowledgements

We appreciate the encouragement, support, and technical assistance of Professor A. A. Loedin, Dr Iskak Koinman, Mr M. A. Hasibuan and the Department of Bacteriology staff, National Institute of Health Research and Development, Ministry of Health, Jakarta, Indonesia; Dr E. M. P. Tampubolon DOH, Mr Sadhani and Deddy Djunardi, Department of Health, Pertamina Oil Company, Jakarta and

Pisju, Indonesia; Dr Larry W. Laughlin, Dr Fred P. Paleologo, Dr Narain H. Punjabi, Decy Subekti and Senhawi Sadeli of US NAMRU-2, Jakarta, Indonesia; and Betty Leckey, Naval Medical Research Institute, Bethesda, Maryland, USA.

This study was supported by the World Health Organization fund No. 83141, T6/181/25; Ministry of Health, Indonesia; Pertamina Oil Co., Jakarta, Indonesia; and the Naval Medical Research and Development Command, Navy Department, Work Unit No. 3M161102BS10.AF.428.

References

- Benavente, L., Gotuzzo, E., Guerra, J., Grados, O., Guerra, H. & Bravo, N. (1984). Diagnosis of typhoid fever using a string capsule device. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 78, 404-406.
- Edwards, P. R. & Ewing, W. H. (1972). *Identification of Enterobacteriaceae*, 3rd edition. Minneapolis: Burgess Publishing Co.
- Gilman, R. H., Terminel, M., Levine, M. M., Hernandez-Mendoza, P. & Hornick, R. B. (1975). Relative efficacy of blood, urine, rectal swab, bone-marrow and rose-spot cultures for recovery of *Salmonella typhi* in typhoid fever. *Lancet*, i, 1211-1213.
- Guerra-Caceras, J. G., Gotuzzo-Herencia, E., Corby-Dagnino, E., Miro-Quezada, M. & Carrillo-Parodi, C. (1979). Diagnostic value of bone-marrow culture in typhoid fever. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 73, 680-683.
- Hoffman, S. L., Punjabi, N. H., Rockhill, R. C., Sutomo, A., Rivai, A. R. & Pulungaih, S. P. (1984). Duodenal string-capsule culture compared with bone-marrow, blood and rectal swab cultures for diagnosing typhoid and paratyphoid fever. *Journal of Infectious Diseases*, 149, 157-161.
- Hoffman, S. L., Edman, D. C., Punjabi, N. H., Lesmana M., Cholid, A., Sundah, S., & Harahap, J. (1986). Bone marrow aspirate culture superior to streptokinase clot culture and 8 ml 1:10 blood-to-broth ratio blood culture for diagnosis of typhoid fever. *American Journal of Tropical Medicine and Hygiene*, 35, 836-839.
- Watson, K. C. (1978). Laboratory and clinical investigations of recovery of *Salmonella typhi* from blood. *Journal of Clinical Microbiology*, 2, 122-126.

Received 3 August 1987; accepted for publication 2 September 1987